

Summary Sheet

Please detach and keep or discard after the exam.

0.1 Notation

- (model) $X \sim \mathbb{P}_\theta$ with values in \mathcal{X} , where $\theta \in \Omega$ is the parameter
- (loss and risk) $L(\theta, d)$ and $R(\theta, \delta) = \mathbb{E}_\theta[L(\theta, \delta(X))]$, where $\delta(X)$ is a statistic
- (average risk) For a prior Λ and estimator δ , we denote by $r(\Lambda, \delta) = \int R(\theta, \delta)\Lambda(d\theta)$, which is the average risk with respect to Λ . We also let δ^Λ denote a Bayes estimator (when one exists) and $r_\Lambda = r(\Lambda, \delta^\Lambda)$ the average risk of that estimator, also called the Bayes risk.
- (maximum risk) For an estimator δ we let $\bar{R}(\delta) = \sup_{\theta \in \Omega} R(\theta, \delta)$, which is its maximum risk.

0.2 Equivariance setting

- We say that a transformation $g : \mathcal{X} \rightarrow \mathcal{X}$ leaves the model invariant if (i) it is one-to-one; (ii) if for any $\theta \in \Omega$, $X \sim \mathbb{P}_\theta$ implies $gX \sim \mathbb{P}_{\theta'}$ for some $\theta' \in \Omega$, which allows one to define $\bar{g} : \Omega \rightarrow \Omega$ that associates θ' to θ ; (iii) \bar{g} is one-to-one (here we assume the model is identifiable).
- We work with a group G of transformations, each leaving the model invariant, and define $\bar{G} = \{\bar{g} : g \in G\}$.
- We work with an estimand $h(\theta)$ that is equivariant with respect to \bar{G} . This allows one to define g^* for each $g \in G$ such that $h(\bar{g}\theta) = g^*h(\theta)$ for all $\theta \in \Omega$.
- We work with a loss function L that is invariant in that $L(\bar{g}\theta, g^*d) = L(\theta, d)$ for all θ , all d , and all g .
- We then focus on estimators that are equivariant in the sense that $\delta(gx) = g^*\delta(x)$ for all x and all g .

0.3 Location model

This model is of the form $X = (X_1, \dots, X_n) \sim f(x - \xi)$ where f is a given density on \mathbb{R}^n and $\xi \in \mathbb{R}$ is unknown. (As usual, $x - \xi$ is understood coordinate-wise.) The transformations of interest are of the form $x \mapsto a + x$ where $a \in \mathbb{R}$. We want to estimate ξ and work with a loss of the form $L(\xi, d) = \rho(d - \xi)$. Let $Y = (Y_1, \dots, Y_{n-1})$ with $Y_i = X_i - X_n$.

Theorem 1. *In the present setting, let δ_0 be any equivariant statistic with finite risk, and suppose we may define $v^*(y) = \arg \min_{v \in \mathbb{R}} \mathbb{E}_0[\rho(\delta_0(X) - v) \mid Y = y]$. Then $\delta^*(x) = \delta_0(x) - v^*(y)$ is MRE.*

Corollary 1. *Under squared error loss, the MRE may be expressed as $\delta^*(x) = \int_{-\infty}^{\infty} v f(x - v) dv / \int_{-\infty}^{\infty} f(x - v) dv$.*

Proposition 1. *Under squared error loss, if an UMVUE exists and is equivariant, then it is MRE.*

0.4 Scale model

This model is of the form $X = (X_1, \dots, X_n) \sim \tau^{-1} f(x/\tau)$ where f is a given density on \mathbb{R}^n and $\tau > 0$ is unknown. The transformations of interest are of the form $x \mapsto bx$ where $b > 0$. We want to estimate τ and work with a loss of the form $L(\xi, d) = \gamma(d/\tau)$. Let $Z = (Z_1, \dots, Z_n)$ with $Z_i = X_i/X_n$ for $i \neq n$ and $Z_n = X_n/|X_n|$.

Theorem 2. *In the present setting, let δ_0 be any equivariant statistic with finite risk, and suppose we may define $w^*(z) = \arg \min_{w > 0} \mathbb{E}_1[\gamma(\delta_0(X)/w) \mid Z = z]$. Then $\delta^*(x) = \delta_0(x)/w^*(z)$ is MRE.*

Corollary 2. *When $L(\tau, d) = (d/\tau - 1)^2$, the MRE may be expressed as $\delta^*(x) = \int_0^\infty w^n f(wx) dw / \int_0^\infty w^{n+1} f(wx) dw$.*

0.5 Location-scale model

This model is of the form $X = (X_1, \dots, X_n) \sim \tau^{-n} f((x - \xi)/\tau)$ where f is a given density on \mathbb{R}^n and $\xi \in \mathbb{R}$ and $\tau > 0$ are both unknown. The transformations of interest are of the form $x \mapsto a + bx$ where $a \in \mathbb{R}$ and $b > 0$. When our goal is to estimate ξ , we work with a loss of the form $L(\xi, \tau; d) = \rho((d - \xi)/\tau)$. When our goal is to estimate τ , we work with a loss of the form $L(\xi, \tau; d) = \gamma(d/\tau)$.

0.6 Bayes estimation

- B1 Under loss $L(\theta, d) = w(\theta)(d - h(\theta))^2$, the Bayes estimator is $\delta^\Lambda(x) = \mathbb{E}[w(\theta)h(\theta) | X = x] / \mathbb{E}[w(\theta) | X = x]$.
- B2 In a Bayesian setting, suppose the loss is strictly convex (in d) and that Q denotes the marginal of X . Then the Bayes estimator is unique if the Bayes risk is finite and, for any measurable set A , $Q(A) = 0$ implies $P_\theta(A) = 0$ for all θ .

0.7 Minimax estimation

- M1 Suppose that Λ is a prior such that $r_\Lambda = \bar{R}(\delta^\Lambda)$. Then δ^Λ is minimax, and uniquely so if it is unique Bayes.
- M2 If an estimator is Bayes for some prior and has constant risk, it is minimax.
- M3 If, for an estimator δ , we can find a sequence of priors (Λ_k) such that $\liminf_k r_{\Lambda_k} \geq \bar{R}(\delta)$, then δ is minimax.
- M4 Consider $\Omega_0 \subset \Omega$. If an estimator is minimax over Ω_0 and achieves its maximum risk at some $\theta \in \Omega_0$, then this estimator is also minimax over Ω .

0.8 Admissibility

- A1 A unique Bayes estimator is admissible.
- A2 (Karlin's theorem) Suppose $X \sim f_\theta$, where $f_\theta(x) = \beta(\theta)e^{\theta T(x)}$ with respect to some underlying measure. Let $\Omega = [\theta_*, \theta^*]$ denote the natural parameter space. Suppose $L(\theta, d) = (d - h(\theta))^2$, where $h(\theta) = \mathbb{E}_\theta(T)$. Then, for $a \geq 0$ and $b \in \mathbb{R}$, a sufficient condition for $\frac{1}{1+a}T + \frac{a}{1+a}b$ to be admissible is that $\int_{\theta_*}^0 K(\theta)d\theta = \infty$ and $\int_0^{\theta^*} K(\theta)d\theta = \infty$, where $K(\theta) = e^{-ba\theta}\beta(\theta)^{-a}$.
- A3 If an estimator has constant risk and is admissible, it is minimax.
- A4 If an estimator is unique minimax, it is admissible.