

Question I. $U_1, \dots, U_n \rightarrow \text{iid unif}[\beta, \beta+1]; \beta \in \mathbb{R}; n \geq 2$

- (a) Explain whether this is a location family, or a scale family, or perhaps both.
- (b) Show that $(U_{(1)}, U_{(n)})$ is sufficient but not complete.
- (c) Show that $(U_{(1)} + U_{(n)})/2$ is not sufficient
- (d) Show that $\bar{U} - \frac{1}{2}$ is not UMVU for β
- (e) Find the ARE of $\bar{U} - \frac{1}{2}$ ($= \hat{\beta}_1$ say) to median $\{X_i\} - \frac{1}{2}$ ($= \hat{\beta}_2$ say) as estimators of β in this problem.
- (f) Derive the MLE $\hat{\beta}$ for this problem. What is its consistency rate? Support your answer with a rough argument, and if the rate is not the usual \sqrt{n} -rate, what regularity conditions are violated to make the usual theorem's conclusion invalid?

Question II. A toy ^{testing} problem: $\Theta = \{0, 1\}$, Z is data taking values in a 3-point set $\{1, 2, 3\}$.

Null $H_0: \theta = 0; Z \sim f_0(z)$
 Alternative $H_1: \theta = 1; Z \sim f_1(z)$ } discrete distributions, tabulated below.

	$z=1$	$z=2$	$z=3$
$f_0(z)$	$\frac{1}{4}$	0	$\frac{3}{4}$
$f_1(z)$	$\frac{1}{2}$	$\frac{1}{2}$	0

(a) Find the most powerful test of $H_0: \theta = 1$ vs. $H_1: \theta = 2$, at level $\alpha = 0.25$, and find its power.

(b) Why does a most powerful test (simple vs. simple) always have power at least as large as its significance level?

(c) Find a test based on the likelihood ratio (as in (a)) with level $\alpha = 0.3$. It will be a randomized test. Explain how you perform it using Z and a coin.

(d) Again another distribution f_2 to the alternative hypothesis (still supported in $\{1, 2, 3\}$) so your test in (a) is still UMP against $H_2 = \{f_1, f_2\}$.

(e) Again a third distribution f_3 so your test in (a) is no longer UMP against $\{f_1, f_2, f_3\}$.

(f) A Bayesian sets a prior π on Θ with $\pi(0) = 0.4$, $\pi(1) = 0.6$.

Derive the Bayes rule for estimating θ with squared-error loss, and compute the Bayes risk of the Bayes rule.

(g) Is the prior in (f) least-favorable? What is?

Question III. (On skewness estimation)

Recall that when $X_1, \dots, X_n \sim \text{iid}(\mu, \sigma^2)$ ($E|X_i|^3 < \infty$) the skewness is defined by $\tau = E[X_i - \mu]^3 / \sigma^3$. (τ is a location- and scale-invariant measure of asymmetry, and its estimation is useful in fitting "Edgeworth corrections" to the CLT approximation.)

The usual estimator under nonparametric assumptions is

$$\hat{\tau} = \hat{\tau}_n = \frac{\frac{1}{n} \sum (x_i - \bar{x})^3}{\left(\frac{1}{n} \sum (x_i - \bar{x})^2\right)^{3/2}} = \frac{\hat{\mu}_3}{\hat{\sigma}_3^3} \text{ say.}$$

- (a) How do we know this is not UMVUE under normality assumptions?
- (b) Give an example of a discrete bivariate distribution, supported on the smallest set you can think of, where the one coordinate r.v. is either independent of, or uncorrelated with, the other, but not both.
- (c) Show that under normality the numerator $\hat{\mu}_3$ and denominator $\hat{\sigma}_3^3$ in $\hat{\tau}$ above have the property in (b) (although of course this is not a discrete situation).
- (d) How do we know readily that under normal sampling $\sqrt{n}(\hat{\sigma}^2 - \sigma^2) \xrightarrow{d} N(0, 2\sigma^4)$? (There are several possible answers.)
- (e) Find a similar weak convergence result for $\hat{\mu}_3$. (The binomial theorem and Slutsky's theorem may come in handy. Also,

you may assume that $E Z^6 = 15$ when $Z \sim \mathcal{N}(0, 1)$.

(f) In class we mentioned without proof a bivariate form of the delta method. Roughly, if $g(\theta_1, \theta_2)$ is nice, and

$$\sqrt{n} \begin{pmatrix} \hat{\theta}_1 \\ \hat{\theta}_2 \end{pmatrix} - \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix} \xrightarrow{d} \mathcal{N}_2 \left(\mathbf{0}, \begin{pmatrix} \sigma_{11}^2 & \sigma_{12} \\ \sigma_{12} & \sigma_{22}^2 \end{pmatrix} \right), \text{ then}$$

$$\sqrt{n} (g(\hat{\theta}_1, \hat{\theta}_2) - g(\theta_1, \theta_2)) \xrightarrow{d} \mathcal{N} \left(0, \sigma_1^2 \left(\frac{\partial g}{\partial \theta_1}(\theta_1, \theta_2) \right)^2 + \sigma_2^2 \left(\frac{\partial g}{\partial \theta_2}(\theta_1, \theta_2) \right)^2 + 2\sigma_{12} \frac{\partial g}{\partial \theta_1}(\theta_1, \theta_2) \cdot \frac{\partial g}{\partial \theta_2}(\theta_1, \theta_2) \right)$$

Using this formula, and simplifying it by assuming $\sigma_{12} = 0$ (because of what you found in (c)), show that, if in fact your original data were from $\mathcal{N}(\mu, \sigma^2)$, $\sqrt{n} \hat{t}_n \xrightarrow{d} \mathcal{N}(0, 6\sigma^4)$.